환경 및 응용기상 분과 [P-084]

Analysis of Information Content of XCO₂ from Remote Sensing for Estimating Surface CO₂ Fluxes

<u>강지순</u>¹, 김원수¹, 명훈주¹, 신설은²

¹한국과학기술정보연구원 ²한국표준과학연구원

Spatial and temporal variation of surface CO_2 fluxes can be estimated by a top-down approach that utilizes both atmospheric CO_2 observations and atmospheric transport model. Even though the transport model is used to link the relationship between surface CO_2 fluxes and atmospheric CO_2 distribution, transport error due to imperfection of the numerical model has been an obstacle to the accuracy of top-down approach (Enting, 2002). One more limitation of topdown approach is the lack of observations. There are few ground-based observations. While satellite observation data cover the Earth fairly well, the information is column-mixed, XCO_2 , according to flat vertical averaging kernels of the sensors. That is, we do not have enough observations to resolve near-surface CO_2 distribution with the issue of transport errors. Therefore, in this study, we would like to figure out how much information has been blurred by column-mixed observations and explore the way to effectively extract near-surface distribution of atmospheric CO_2 from XCO_2 data with a help of numerical model data under the OSSEs with SPEEDY-C. We also tried to generate near-surface atmospheric CO_2 data at the satellite observation space using neural network model. With the OSSEs, we also plan to consider the impact of transport error as a further study.

Key words: Surface CO₂ fluxes, SPEEDY-C, XCO₂, Transport error, Neural Network model

※ 이 연구는 국가 플래그십 초고성능컴퓨터 인프라 구축 및 서비스 사업(과제번호: (KISTI)K24L2M1C1-01, (NTIS)2710006789)의 지원으로 수행되었습니다.